

REALITY CONDITIONS FOR ASHTEKAR VARIABLES AS DIRAC CONSTRAINTS

Hugo A. Morales–Técotl^{1,3*}, Luis F. Urrutia^{1,2} and J. David Vergara^{2†}

¹ Departamento de Física
Universidad Autónoma Metropolitana–Iztapalapa
Apartado Postal 55–534, 09340 México D.F., México

² Instituto de Ciencias Nucleares
Universidad Nacional Autónoma de México
Apartado Postal 70-543, 04510 México D.F., México

³ International Centre for Theoretical Physics,
P.O. Box 586, 34100 Trieste, Italy.

May 1996

Abstract

We show that the reality conditions to be imposed on Ashtekar variables to recover real gravity can be implemented as second class constraints *à la* Dirac. Thus, counting gravitational degrees of freedom follows accordingly. Some constraints of the real theory turn out to be non-polynomial, regardless of the form, polynomial or non-polynomial, taken for the reality conditions. We comment upon the compatibility of our approach with the recently proposed Wick transform point of view, as well as on some alternatives for dealing with such second class constraints.

PACS: 04.60, 04.60.D

*hugo@xanum.uam.mx

†vergara@nuclecu.unam.mx

The formulation of general relativity in terms of the Ashtekar variables: a densitized inverse dreibein $\tilde{e}^{ai}$ together with a complex $so(3)$ connection A_{ai} [1], has opened up a broad new avenue of research [2]. Reexpressing general relativity as a theory of a connection has made it possible to incorporate methods which have been very useful and successful in the description of Yang-Mills theories. A most notable example of this interbreeding is the recent application of the loop-space methods to the quantization of general relativity [3]. Another important advantage of the new variables is that the constraints characterizing general relativity turn out to be polynomial in these phase space variables. This simplicity, however, gets afflicted by the complex character of the new variables: the use of reality conditions is compulsory and the pursue of the Ashtekar program deviates somewhat from the more conventional approaches. In fact, the original Ashtekar program was formulated as an alternative route for classical general relativity and its canonical quantization, as compared to the standard ADM approach, for example. In the classical theory, general relativity is considered to be embedded in a larger, complex theory. The restriction to the usual Einstein theory is subsequently made imposing by hand some reality conditions. In the quantum theory, one first ignores the reality conditions, solves the quantum constraints of the complex theory and finally imposes the reality conditions as constraints upon the admissible inner products by requiring that the real classical observables become self-adjoint operators. The use of the reality conditions as a way of selecting an inner product has worked well in the case of some models, but the same has not happened for full gravity so far [4].

The aim of the present work is to reanalyze classical Ashtekar complex gravity

plus the reality conditions in the light of the canonical Dirac approach to constrained systems. In particular, we want to elucidate the status of the reality conditions viewed as Dirac constraints. A naive analysis of the standard reality conditions: q^{ab} real and $\dot{q}^{ab}$ real, as second class constraints, is misleading because the count of the number of degrees of freedom in the configuration space turns out to be -2 per spatial point, which is obviously wrong! A way out to this situation has been recently proposed in Ref. [5], outside the Dirac framework, however, following a previous result in Ref. [6].

In this work we start with reality conditions, also imposed by hand, but considering them as primary constraints in the Ashtekar self-dual action. Then, we systematically apply the Dirac procedure, showing that these reality conditions end up by being second-class constraints and recovering the standard Hamiltonian formulation of general relativity in its Palatini canonical form [1]. In this way, we show that it is possible to extract a satisfactory canonical description of general relativity starting from the complex theory. In fact, we prove that the original complexified phase-space, both in the dreibein and in the connection, together with the self-dual Ashtekar action plus the reality conditions, lead to the standard phase space of real Einstein gravity, via the Dirac method.

A closely related issue is that of the polynomiality of the remaining first-class constraints, in terms of the canonical variables. It has been shown in Ref. [7] that the elimination of the associated second-class constraints in the complex Palatini action with self-dual connection [8, 9], can be done without destroying the polynomial character of the remaining first-class constraints, as opposed to the real Palatini

case. Nevertheless, this nice property holds only before imposing the required reality conditions. The non-polynomial choice for such reality conditions made in Ref. [7] is briefly discussed there only in relation with the inner product of the quantum theory and no attempt to interpret them as Dirac constraints is made. We consider explicitly a non-polynomial form of the reality conditions and only describe at the end the pertinent changes produced by a polynomial choice. The canonical analysis leads in both cases to a formulation of real general relativity in terms of non-polynomial constraints. This feature seems to be compatible with the recent Wick transform method proposed by Thiemann [10] and Ashtekar [11], that makes non-polynomiality a ‘controllable’ issue. Also, this result seems to be consistent with the point of view taken in Ref. [12], where it is shown that among a subset of the canonical transformations leading to Ashtekar-type variables in the case of Lorentzian signature, only the complex case leads to a polynomial Hamiltonian constraint. Thus, real Ashtekar-type variables can be used *ab initio* provided one is willing to accept a more complicated Hamiltonian constraint.

Let us proceed now to analyze the canonical form of the self-dual action [8, 9]

$$S = \int dt d^3x \left\{ -i\tilde{e}^{ai}\dot{A}_{ai} - N\mathcal{S} - N^a\mathcal{V}_a - N^i\mathcal{G}_i \right\}, \quad (1)$$

where

$$\mathcal{S} := \epsilon_{ijk}\tilde{e}^{ai}\tilde{e}^{bj}F_{ab}{}^k, \quad \mathcal{V}_a := \tilde{e}^b{}_j F_{ab}{}^j, \quad \mathcal{G}_i := \mathcal{D}_a\tilde{e}^a{}_i, \quad (2)$$

are the constraints of the theory and N, N^a, N^i are Lagrange multipliers. The *complex* canonical variables are: i) $\tilde{e}^{ai} := e e^{ai}$, with e^{ai} being the triad ($e^{ai}e^b{}_i := q^{ab}$, q^{ab} is the spatial three-metric), and $a, b, \dots = 1, 2, 3$ are spatial indices, whereas $i, j, \dots =$

1, 2, 3 are $\text{so}(3)$ internal indices. Also $e := \det e_{bj}$ with e_{bj} being the inverse of e^{ai} . ii) A_{ai} is the three-dimensional projection of the selfdual connection [1] with associated covariant derivative $\mathcal{D}_a \lambda_i = \partial_a \lambda_i + \epsilon_{ijk} A_a^j \lambda^k$ and $F_{ab}^i := \partial_a A_b^i - \partial_b A_a^i + \epsilon^i_{jk} A_a^j A_b^k$ is the corresponding curvature. The above complex pair of canonically conjugated variables satisfies

$$\{\tilde{e}^{ai}(x), A_{bj}(y)\} = i \delta_b^a \delta_j^i \delta^3(x, y). \quad (3)$$

With these Poisson brackets, the complex constraints (2) are first class, i.e. for any two of them, say $\mathcal{Q}$ and $\mathcal{R}$, we have

$$\{\mathcal{Q}, \mathcal{R}\}_{A, \tilde{e}} \approx 0. \quad (4)$$

Let us remind the reader of some basic features of the Dirac method (cf [13]), which will be used in what follows. Consider a dynamical system with real canonical variables q^i, p_i $i = 1, \dots, N$ together with a complete set of real constraints $G_a(q, p)$, $a = 1, \dots, m$. The function $F(q, p)$ is called a first-class function provided $\{F, G_a\} = U_a^b(q, p) G_b, \forall a$. If this equality does not hold, $F(q, p)$ is called a second class function. Hereby, the complete set of constraints gets classified into first class constraints, Φ_α , $\alpha = 1, \dots, n \leq m$ and second class constraints, χ_A , $A = n + 1, \dots, m$. The second class set satisfies $\{\chi_A, \chi_B\} = C_{AB}$, with $\det C_{AB} \neq 0$. The number of physical (configuration) degrees of freedom is given by:

$$\frac{1}{2} [\dim(\text{phase space}) - 2(\# \text{ of 1st class constraints}) - (\# \text{ of 2nd class constraints})].$$

In this way, the count of real degrees of freedom corresponding to the action (1) leads to $\frac{1}{2}[2 \times 2 \times 9 - 2 \times 2 \times 7] = 4$ per spatial point, i.e. the action (1) describes

complex gravity. Notice that we have included an additional factor of 2 in this count because of the complex character of both the canonical variables and the constraints. Clearly, one has to supplement the theory with reality conditions in order to go back to real gravity.

We shall proceed with the canonical analysis of the action (1), together with the necessary reality conditions, by splitting the complex canonical variables $A_{ai}, \tilde{e}^{bj}$ into real and imaginary parts according to

$$\tilde{e}^{ai} = E^{ai} + i\mathcal{E}^{ai}, \quad A_{bj} = \gamma_{bj} - iK_{bj}. \quad (5)$$

Most importantly, we will consider all of them (36 real variables) as configuration variables to begin with. Observe that this procedure dispenses us from dealing with the complex conjugate of the Ashtekar variables in the formulation of the reality conditions. This is very convenient because their Poisson brackets would not even be defined in the theory.

The reality conditions we start with are

$$\psi^{ai} := \mathcal{E}^{ai} = 0, \quad \chi_{ai} := \gamma_{ai} - f_{ai}(E) = 0, \quad (6)$$

which are subsequently taken as primary constraints ψ^{ai}, χ_{bj} , supplementing the action (1). We can see that the constraint ψ^{ai} enforces the initially complex triad $\tilde{e}^{ai}$ to be real, and hence the corresponding three-metric. Based upon the compatibility condition between a real torsion-free connection $\overset{0}{\gamma}_{ai}$ and a real densitized triad E^{bj}

$$D_a E^{bj} = \partial_a E^{bj} + \Gamma_{ac}{}^b E^{cj} + \epsilon^j{}_{kl} \overset{0}{\gamma}_a{}^k E^{bl} - \Gamma_{da}{}^d E^{bj} = 0, \quad (7)$$

where $\Gamma_{ab}{}^c$ is the Levi-Civita connection defined by the metric $h^{ab} := E^{ai} E^b{}_i$, we

choose the form of f_{ai} as

$$f_{ai} = \frac{1}{2}[E_{ai}E_c^j\epsilon_{jrs} - 2E_{aj}E_c^j\epsilon_{irs}]E^{dr}\partial_d E^{cs}. \quad (8)$$

In this way we guarantee that γ_{ai} is the real torsion-free connection compatible with E^{bj} , i.e. $\gamma_{ai} = \gamma_{ai}^0$. Since E_{ai} is the inverse of E^{bj} , (8) is non-polynomial in E^{bj} and so it is χ_{ai} .

The definition of the canonical momenta, denoted generically by Π , in the action (1) leads to the following constraints

$$\phi_{\mathcal{E}ai} = \Pi_{\mathcal{E}ai}, \quad \phi_{\gamma}^{ai} = \Pi_{\gamma}^{ai} + iE^{ai}, \quad \phi_K^{ai} = \Pi_K^{ai} + E^{ai}, \quad \phi_{Eai} = \Pi_{Eai}, \quad (9)$$

where we note that Π_{γ}^{ai} is purely imaginary. The corresponding non-zero Poisson brackets are

$$\begin{aligned} \{\mathcal{E}^{ai}(x), \Pi_{\mathcal{E}bj}(y)\} &= \delta_a^b \delta_i^j \delta^{(3)}(x, y), \quad \{E^{ai}(x), \Pi_{Ebj}(y)\} = \delta_a^b \delta_i^j \delta^{(3)}(x, y), \\ \{\gamma_{ai}(x), \Pi_{\gamma}^{bj}(y)\} &= \delta_a^b \delta_i^j \delta^{(3)}(x, y), \quad \{K_{ai}(x), \Pi_K^{bj}(y)\} = \delta_a^b \delta_i^j \delta^{(3)}(x, y). \end{aligned} \quad (10)$$

In this way, the full set of primary constraints is formed by (2), (6) and (9), written in terms of the real variables (5). Remarkably, the analysis of the phase space could be also performed starting from a real triad $\tilde{e}^{ai}$. In this way, the dimension of the original phase space is reduced, together with the number of required reality conditions. This alternative procedure shows that there is nothing wrong in considering a “mixed phase space”: real $\tilde{e}^{ai}$ and complex A_{bj} , as long as the canonical analysis is consistently carried out from the very beginning. Nevertheless, we choose to deal with both complex configuration variables because we obtain a more symmetrical description in doing so.

The next step is to analyze the time evolution of the primary constraints, together with the classification of the resulting set of constraints. It turns out to be appropriate to redefine

$$\phi_{Eai} \rightarrow \phi'_{Eai} = \phi_{Eai} + \alpha_{aibj} \phi_\gamma^{bj} + \beta_{ai}^{bj} \chi_{bj} + \eta_{aibj} \phi_K^{bj}, \quad (11)$$

with

$$\begin{aligned} \alpha_{aibj}(x, y) &= \frac{\delta f_{bj}(y)}{\delta E^{ai}(x)}, \quad \beta_{ai}^{bj}(x, y) = i \delta_a^b \delta_i^j \delta^{(3)}(x, y), \\ \eta_{aibj}(x, y) - \eta_{bjai}(y, x) &= i \left(\frac{\delta f_{ai}(x)}{\delta E^{bj}(y)} - \frac{\delta f_{bj}(y)}{\delta E^{ai}(x)} \right), \end{aligned} \quad (12)$$

where $\frac{\delta f(y)}{\delta E(x)}$ denotes the corresponding functional derivative. Our notation, introduced in Eq.(11), is that the summation over repeated indices involves also an integral over the three dimensional space. In this way, for example $\phi_A = \alpha_{AB} \chi^B$ stands for $\phi_A(\mathbf{x}, t) = \int d^3\mathbf{y} \alpha_{AB}(\mathbf{x}, \mathbf{y}) \chi^B(\mathbf{y}, t)$.

The resulting Poisson brackets matrix for the subset of constraints (6), (9) and (11) is

	$\phi_{\mathcal{E}}$	ψ	ϕ_γ	χ	ϕ_K	ϕ'_E	
$\phi_{\mathcal{E}}$	0	$-\delta_a^b \delta_i^j \delta^{(3)}$	0	0	0	0	
ψ	$\delta_b^a \delta_j^i \delta^{(3)}$	0	0	0	0	0	
ϕ_γ	0	0	0	$-\delta_b^a \delta_j^i \delta^{(3)}$	0	0	
χ	0	0	$\delta_a^b \delta_i^j \delta^{(3)}$	0	0	0	
ϕ_K	0	0	0	0	0	$\delta_a^b \delta_j^i \delta^{(3)}$	
ϕ'_E	0	0	0	0	$-\delta_b^a \delta_i^j \delta^{(3)}$	0	, (13)

where $\delta^{(3)} = \delta^{(3)}(x, y)$. From the above matrix we conclude that this subset, which includes the reality conditions, is second class. Since the original constraints (2)

generate the gauge symmetries of the system, we would like to keep them as first-class constraints. To see whether this is tenable, we start by searching a redefinition of them so that they have zero Poisson brackets with the second-class set. Notice that the constraints (2) only depend on the configuration variables and not on the momenta. For any of them, say $\mathcal{R}$, we construct

$$\mathcal{R}' = \mathcal{R} + \{\phi_{\mathcal{E}bj}, \mathcal{R}\}\psi^{bj} + \{\phi_{\gamma}^{bj}, \mathcal{R}\}\chi_{bj} + \{\phi'_{Ebj}, \mathcal{R}\}\phi_K^{bj} - \{\phi_K^{bj}, \mathcal{R}\}\phi'_{Ebj}, \quad (14)$$

which we can verify to fulfill the above requirement. It remains now to calculate the Poisson brackets among the constraints of the set $\mathcal{S}', \mathcal{V}'_a, \mathcal{G}'_i$. Let us consider $\{\mathcal{Q}', \mathcal{R}'\}$, i.e. any two of them

$$\begin{aligned} \{\mathcal{Q}', \mathcal{R}'\} \approx \{\mathcal{Q}', \mathcal{R}\} &= \{\phi_{\mathcal{E}ai}, \mathcal{Q}\}\{\psi^{ai}, \mathcal{R}\} + \{\phi_{\gamma}^{ai}, \mathcal{Q}\}\{\chi_{ai}, \mathcal{R}\} \\ &+ \{\phi'_{Eai}, \mathcal{Q}\}\{\phi_K^{ai}, \mathcal{R}\} - \{\phi'_{Eai}, \mathcal{R}\}\{\phi_K^{ai}, \mathcal{Q}\}. \end{aligned} \quad (15)$$

The first two terms yield zero because the constraints ψ^{ai}, χ_{ai} have no dependence on the momenta. After substituting the expression (11) for ϕ'_{Eai} we are left with

$$\begin{aligned} \{\mathcal{Q}', \mathcal{R}'\} \approx \{\phi_K^{ai}, \mathcal{R}\}\{\phi_{Eai}, \mathcal{Q}\} &- \{\phi_K^{ai}, \mathcal{Q}\}\{\phi_{Eai}, \mathcal{R}\} \\ &+ (\eta_{aibj} - \eta_{bjai})\{\phi_K^{ai}, \mathcal{R}\}\{\phi_K^{bj}, \mathcal{Q}\} \\ &+ \alpha_{aibj} \left[\{\phi_K^{ai}, \mathcal{R}\}\{\phi_{\gamma}^{bj}, \mathcal{Q}\} - \{\phi_K^{ai}, \mathcal{Q}\}\{\phi_{\gamma}^{bj}, \mathcal{R}\} \right]. \end{aligned} \quad (16)$$

A straightforward calculation based on the splitting (5) for the first line in (16) leads to

$$\begin{aligned} \{\phi_K^{ai}, \mathcal{R}\}\{\phi_{Eai}, \mathcal{Q}\} - \{\phi_K^{ai}, \mathcal{Q}\}\{\phi_{Eai}, \mathcal{R}\} &= -i \left(\frac{\delta \mathcal{R}}{\delta A_{ai}} \frac{\delta \mathcal{Q}}{\delta \tilde{e}^{ai}} - \frac{\delta \mathcal{Q}}{\delta A_{ai}} \frac{\delta \mathcal{R}}{\delta \tilde{e}^{ai}} \right) \\ &= -i \{\mathcal{R}, \mathcal{Q}\}_{A, \tilde{e}} \approx 0, \end{aligned} \quad (17)$$

according to the first-class property (4) of $\mathcal{R}$, $\mathcal{Q}$ in the original complex phase space. Besides, taking into account the corresponding coefficients (12) the second line of (16) reduces to

$$i \left(\frac{\delta f_{bj}}{\delta E^{ai}} - \frac{\delta f_{ai}}{\delta E^{bj}} \right) \frac{\delta \mathcal{R}}{\delta A_{ai}} \frac{\delta \mathcal{Q}}{\delta A_{bj}},$$

while the third line is just the negative of this same expression. We have thus shown that the Poisson brackets among any pair of the set of constraints $\mathcal{S}'$, $\mathcal{V}'_a$, $\mathcal{G}'_i$ is weakly zero.

Summarizing, our system is conveniently described by the following set of primary constraints: $\phi_{\mathcal{E}ai}$, ϕ_{γ}^{ai} , ϕ_K^{ai} , ϕ'_{Eai} , ψ^{ai} , χ_{ai} , $\mathcal{S}'$, $\mathcal{V}'_a$, $\mathcal{G}'_i$. Next, we impose the conservation in time of this whole set, using the total hamiltonian density

$$\begin{aligned} \mathcal{H}_{\text{Total}}(\mathbf{x}) \approx & \mu_{\mathcal{E}}^{ai} \phi_{\mathcal{E}ai} + \mu_{\gamma ai} \phi_{\gamma}^{ai} + \mu_{Kai} \phi_K^{ai} + \mu_E^{ai} \phi'_{Eai} + \lambda_{ai} \psi^{ai} + \omega^{ai} \chi_{ai} \\ & + N \mathcal{S}' + N^a \mathcal{V}'_a + N^i \mathcal{G}'_i, \end{aligned} \quad (18)$$

where no integration over the three-space is involved. The properties of the constraints $\mathcal{S}'$, $\mathcal{V}'_a$, $\mathcal{G}'_i$, together with Eqs. (14) and (17), guarantee that their Poisson brackets with $H_{\text{Total}} = \int d^3\mathbf{x} \mathcal{H}_{\text{Total}}(\mathbf{x})$ is weakly zero. Let us consider now the remaining constraints. By virtue of the Poisson brackets matrix (13) together with the definition (14), the calculation of the corresponding time evolution fixes each of the Lagrange multipliers $\mu_{\mathcal{E}}^{ai}$, $\mu_{\gamma ai}$, μ_{Kai} , μ_E^{ai} , λ_{ai} , ω^{ai} equal to zero. We conclude that there are no secondary constraints and that the set $\mathcal{S}'$, $\mathcal{V}'_a$, $\mathcal{G}'_i$ is first class while the set $\phi_{\mathcal{E}ai}$, ϕ_{γ}^{ai} , ϕ_K^{ai} , ϕ'_{Eai} , ψ^{ai} , χ_{ai} , which includes the reality conditions, is second class.

To make sure that ultimately we are dealing with real gravity we still have to get

the correct number of degrees of freedom, using the Dirac prescription. Also we must recover the specific form of the constraints characterizing real gravity. To do so we start by examining the complex character of each constraint. By simple inspection of (6) and (9) we realize that each of the second class constraints is either purely imaginary or purely real. To proceed with the count we impose them strongly. This leaves us with $2 \times 9 \times 4 - 6 \times 9 = 18$ as the dimension of partially reduced real phase space. The first class constraints remain first class in the resulting Dirac brackets. As shown below, this first class set contains seven constraints each of which is either purely real or purely imaginary. In this way the final count yields $\frac{1}{2} [18 - 2 \times 7] = 2$ real configuration degrees of freedom per spatial point.

Next we verify our statement regarding the reality properties of the first class constraints and get their explicit form. Consider the Gauss constraint in the partially reduced phase space, to begin with. It becomes

$$\mathcal{G}^i \approx \mathcal{D}_a E^{ai} = \partial_a E^{ai} + \epsilon^i_{jk} \gamma_a^j E^{ak} - i \epsilon^i_{jk} K_a^j E^{ak} \approx -i \epsilon^i_{jk} K_a^j E^{ak}, \quad (19)$$

where we have used the compatibility condition (7). Hence $\mathcal{G}^i$ is purely imaginary. For the vector constraint we have

$$\begin{aligned} \mathcal{V}_a \approx E^b_i F_{ab}^i &= E^b_i R_{ab}^i - \epsilon^i_{jk} E^b_i K_b^j K_{aj} - i D_a (K_b^i E^b_i) + i D_b (K_a^i E^b_i) \\ &\approx i \left[D_b (K_a^i E^b_i - \delta_a^b K_c^i E^c_i) \right] \end{aligned} \quad (20)$$

with $R_{ab}^i = 2\partial_{[a}\gamma_{b]}^i + \epsilon^i_{jk}\gamma_a^j\gamma_b^k$. The last line follows from the term containing R being zero due to the Bianchi identity and the term quadratic in K being weakly zero because of (19). We see then that the vector constraint is purely imaginary in

the reduced phase space. At last, the scalar constraint can be rewritten as

$$\mathcal{S} \approx \epsilon_{ijk} E^{ai} E^{bj} F_{ab}{}^k \approx -qR + (E^{al} E^{bm} - E^{bl} E^{am}) K_{bl} K_{am}, \quad (21)$$

where use has been made of the Gauss law (19) together with the compatibility condition (7) to arrive at the final result. Hereby the scalar constraint is purely real. In each case we have recovered the real first class constraint corresponding to the Palatini canonical formulation of real general relativity, as expected.

Had we started from a real triad together with a complex connection, the dimension of the original phase space would have been 54. The definition of the momenta would have produced 27 second class constraints to which we should add 9 reality conditions. Taking into account the remaining 7 first class constraints, the count of the degrees of freedom turns out to be correct one (i.e. two per spatial point), since any of the above constraints is either purely real or purely imaginary.

The analysis performed above was developed using the non-polynomial reality conditions (6), (8) and it follows that the final structure of the complete set of constraints is non-polynomial. One could think that changing the reality conditions to a polynomial form would solve this issue. Unfortunately, the answer is in the negative. For instance, choosing

$$\tilde{\chi}_m{}^a = \epsilon_{mli} E^{bl} \partial_b E^{ai} + 2\gamma_{b[m} E^a{}_{l]} E^{bl}, \quad (22)$$

instead of the second equation in (6), yields a phase space dependent matrix (13) and non-polynomial first class constraints (14).

Our results seem to be compatible with the recently proposed Wick transform relating real Riemannian and real Lorentzian canonical gravity, which automatically

implements the reality conditions [10, 11]. This happens in two respects. First, non-polynomiality always shows up at some stage. In our case we face non-polynomiality when applying the Dirac method, even if we start from polynomial reality conditions. On the other hand, in the Wick transform framework non-polynomiality is manifest in the transform itself. In the latter case, this issue seems to be under control, although a definite answer has not been attained yet [11]. Second, we were able to carry over our analysis to the Riemannian case, showing that the Wick transform takes the full set of Riemannian constraints to the Lorentzian ones, *in* complexified phase space. This reinforces the idea that the natural home for such a transform is the complexified phase space, as opposed to a space-time interpretation which seems improbable [11].

We have shown that the reality conditions can be taken as second class constraints in the Dirac formalism. The standard treatment calls for the elimination of them in terms of Dirac brackets. Nevertheless, this procedure would destroy the advantages of the Ashtekar variables. To surmount this difficulty, we should look for new directions if we want to keep the canonical framework. We can mention two possible strategies along these lines:

i) One can transform the set of second class constraints to a first class set by adding new variables [13]. This transformation is rather straightforward because the Poisson brackets matrix of the second class constraints has the very simple structure given in (13). For example one can transform the canonical pair ψ , $\phi_{\mathcal{E}}$, Eqs. (6, 9) in the following form

$$\tilde{\phi}_{\mathcal{E}ai} := \Pi_{\mathcal{E}ai} + \tilde{\Pi}_{\mathcal{E}ai}, \quad \tilde{\psi}^{ai} := \mathcal{E}^{ai} - \tilde{\mathcal{E}}^{ai}, \quad (23)$$

where the new variables $\tilde{\Pi}_{\mathcal{E}ai}$ and $\tilde{\mathcal{E}}^{ai}$ are canonically conjugated, *i.e.* $\{\tilde{\Pi}_{\mathcal{E}ai}, \tilde{\mathcal{E}}^{bj}\} = \delta_a^b \delta_i^j \delta^{(3)}$. The new constraints $\tilde{\phi}_{\mathcal{E}ai}$ and $\tilde{\psi}^{ai}$ have zero Poisson brackets with the whole set of constraints, then they are first class. Following the same procedure one can transform the complete set of second class constraints into a first class set. Since now all the constraints are first class, they would define the physical states once they are promoted to operators.

ii) Another possibility could be to assume that one of the constraints in each canonical pair of second class constraints can be considered as first class, while the companion is taken to be a gauge fixing condition. This procedure would not require the addition of extra variables but its consistency should be carefully studied.

Both strategies should directly define the physical states of real gravity, having the correct number of two configuration degrees of freedom per spatial point. Regarding the issue of either non-polynomiality or, perhaps, high order polynomiality, further work has to be done along these lines before a conclusion can be achieved. It should be clear that the present results do not preclude accounting for the reality conditions via either the quantum inner product [1] or the Wick transform [10, 11].

Finally, we expect that the present canonical analysis will remain valid for the cases which include matter couplings as well as space-times with boundaries.

Acknowledgements

We thank A. Ashtekar for enlightening discussions regarding this work, as well as R. Gambini for rearousing our interest in this problem. We also acknowledge useful conversations with F. Barbero, G. Immirzi and J. Stephany. The work of HAMT and

LFU has been supported in part by the grant CONACyT # 3544-E9311. JDV and LFU also acknowledge the support from grant UNAM-DGAPA-IN-100694. HAMT would like to thank D.W. Sciama and ICTP-Trieste for supporting a visit upon which the present work was completed and SISSA-Trieste for the warm hospitality extended to him.

References

- [1] Ashtekar A 1986 *Phys. Rev. Letts.* **57** 2244; 1987 *Phys. Rev.* **D36** 1587; 1991 *Lectures on Non-Perturbative Canonical Gravity (Notes prepared in collaboration with R S Tate)* (Singapore: World Scientific) and references therein.
- [2] Bibliography of publications related to Classical and Quantum Gravity in terms of the Ashtekar Variables, updated by T. Schilling GRQC 9409031.
- [3] Rovelli C and Smolin L 1988 *Phys. Rev. Letts.* **61** 1155; 1990 *Nucl. Phys.* **B331** 80; Gambini R 1991 *Phys. Lett.* **B235** 180.
- [4] Kuchař K 1993 *Canonical quantum gravity* preprint GRQC 9304012; Ashtekar A, Lewandowski J, Marolf D, Murão J and Thiemann T 1995 *J. Math. Phys.* **36** 6456.
- [5] Immirzi G 1993 *Class. Quant. Grav.* **10** 2347; Yoneda G and Shinkai H 1996 *Class. Quant. Grav.* **13** 783.
- [6] Ashtekar A, Romano J and Tate R S 1989 *Phys. Rev.* **D40** 2572.
- [7] Ashtekar A, Balachandran A P and Jo S 1989 *Int. J. Mod. Phys.* **A4** 1493.

- [8] Samuel J 1987 *Pramana J. Phys.* **L429** 28.
- [9] Jacobson T and Smolin L 1987 *Phys. Lett.* **B196** 39; 1988 *Class. Quant. Grav.* **5** 583.
- [10] Thiemann T 1995 *Class. Quant. Grav.* **13** 1383.
- [11] Ashtekar A 1995 *A Generalized Wick Transform for Gravity* preprint GRQC 9511083.
- [12] Barbero J F 1995 *Phys. Rev.* **D51** 5498; 1995 *Class. Quant. Grav.* **12** L5.
- [13] Henneaux M and Teitelboim C 1992 *Quantization of Gauge Systems* (Princeton University Press, New Jersey).